

Wound infection made easy



Introduction

Wound healing is a complex, multifaceted process influenced by intrinsic and extrinsic factors, some of which can be controlled. When healing stalls and certain signs and symptoms are present, the wound may be critically colonised or infected. There is no single scientific test to definitively diagnose infection; wound infection is diagnosed by clinical assessment of the wound and the whole patient. It is therefore important that clinicians understand how infection develops, how signs and symptoms manifest in various aetiologies, and how and when to initiate use of topical antimicrobials and systemic antibiotics in wounds clinically diagnosed as infected.

Authors: Swanson T, Grothier L, Schultz G. Full author details can be found on p5.

Wound healing and infection

Wound healing is a complex process that follows a progressive three-step sequence: inflammatory, proliferative and remodelling. These phases can overlap¹, and each stage's duration will be influenced by a variety of factors. When a wound fails to progress to healing or respond to treatment over the expected healing time frame (depending on the patient and wound type), it usually stalls in the inflammatory phase. This non-healing phase, called 'chronicity', has various causative factors². Wound infection often causes chronicity and, therefore, it is important to understand why wound infection occurs, and how to identify and manage it.

Assessing the risk of wound infection

Patient assessment should include the general medical condition and risk factors for wound infection. Age (e.g. neonatal and elderly patients), certain chronic medical conditions (e.g. diabetes, pulmonary disease, vascular disease), medications (e.g. oncology drugs, anti-platelet drugs, glucocorticoid steroids) and lifestyle factors (e.g. smoking, alcohol consumption) put patients with wounds at greater risk for the development of infection³.

Recognising colonisation and infection

There is no single test to definitively diagnose infection; wound infection is diagnosed by clinical assessment of the wound and the whole patient. Swabbing a wound helps determine

sensitivities or resistance to empirical treatment⁴. However, wound swab culture results may be misleading, as clinical microbiology laboratories use methods that select for planktonic bacteria or are not always suitable for culture of anaerobic species. Thus a wound culture might not capture bacteria protected within a biofilm, nor will it detect biofilm, so the result is often inconclusive^{5,6}.

If a swab is needed to determine whether bacteria are present in deeper wound bed tissues, the wound should first be irrigated with normal saline, to remove surface debris and avoid detection of only surface contaminants. The swab should be performed according to solid clinical rationale, per local protocol (Table 1).

Clinical judgement is therefore needed to interpret signs and symptoms. The classic signs and symptoms of wound infection include inflammation, new or increasing pain, local heat, swelling, advancing redness and purulence⁵. However, these indicators are more likely to appear in acute rather than chronic wound infection and, therefore, clinicians must be familiar with the secondary signs of infection. The secondary signs of infection are suggestive of a local wound infection, critical colonisation or high bioburden in a chronic wound¹⁰⁻¹².

It can be particularly difficult to diagnose bioburden levels in wounds with persistent inflammation, such as mixed-aetiology leg ulcers. In such wounds, it is more likely to be critical colonisation or local infection (Figure 1), the key criteria for which include malodour, friable/bleeding tissue, breakdown/increase in size of wound, discolouration, spreading erythema, change in the nature of pain, bridging of the wound, pocketing at the base of the wound, and development of pus/slough or an abscess¹¹. In addition, increased exudate or exudate that has become purulent can be signs that the microbial burden in the wound may be a factor adding to chronicity⁵.

Critical colonisation and local infection can also be difficult to diagnose in diabetic foot ulcers (DFUs), because vascular compromise and neuropathy in the foot often mean that the classic signs of infection are not present¹³. The high morbidity and mortality associated with infection in DFUs mean that early and aggressive treatment — in the presence of even subtle signs of infection — is more appropriate than for wounds of other aetiologies¹⁴ (Table 2).

Role of biofilms

Biofilms will often be present — up to 60% of the time — in wounds with chronicity, and may be responsible for the state of persistent inflammation that makes diagnosing wound infection difficult²¹. Biofilms are dynamic communities of bacteria and fungi

Wound infection made easy



Table 1 When to initiate a wound culture swab⁷⁻⁹

When to swab	Rationale
Initial presentation of symptoms of infection	To determine resistance to empirically commenced antibiotic; to assess virulence and type of wound microbes
Wound(s) not progressing after 2 weeks of treatment	To determine if another causative microbe is active or if antibiotic resistance has occurred
Normal surveillance protocol	Per protocol to screen for certain resistant microbes
Treatment completion (if local protocol advises)	To determine clearance of microbes

Box 1 Potential triggers for use of a topical antimicrobial as prophylaxis³

- Patient history of delayed wound healing/infection
- Gross contamination that presents risk of cross-infection with multidrug-resistant bacteria
- Anatomical location of the wound that increases risk of contamination (e.g. sacrum)
- Wound of 4+ weeks' duration at initial presentation for treatment, with no visible signs of healing, or with signs of continued deterioration
- Evidence of pathologies or activities likely to compromise immunity (e.g. in diabetes with poorly controlled blood glucose, smoking, regular alcohol use beyond recommendations)
- Significantly compromised blood flow where healing is unlikely without vascular intervention
- Odour that affects quality of life




Figure 1. The continuum of wound infection (illustration courtesy G Schultz)

living within a protective self-secreted matrix of sugars and proteins²². They can develop within 2–4 days of initial colonisation, and become very tightly attached to extracellular matrix components or the wound bed, making them difficult to remove by surface irrigation or superficial debridement²³.

Goals of treatment

The wound management plan should be tailored according to the

patient's individual situation. Perform a thorough holistic assessment of the patient's medical history and status, and of the wound and its characteristics to determine the extent to which critical colonisation or local infection is a risk.






Where there is a high risk, but the signs and symptoms of critical colonisation and local infection are absent, the goal of treatment should be to prevent onset of infection. Non-viable and devitalised

tissue provides an opportunity for microbial growth and should therefore be removed from the wound bed and wound edge through irrigation or debridement³. Moisture imbalance should be rectified to optimise a moist wound-healing environment in which exudate is managed and drying out prevented³. In addition, the use of a topical antimicrobial in certain high-risk individuals/wounds (Box 1) can also help prevent an increase in microbial bioburden and the possible onset of infection; the rationale for application of the topical antimicrobial should be clearly documented in the patient's wound management plan and a review date set³.

In wounds exhibiting signs and symptoms of critical colonisation or local infection, the treatment goal must be to reduce the bioburden and/or eradicate potential biofilm in the wound bed. The first step in these wounds is aggressive debridement of slough and the underlying tissue, to disrupt the microbial burden and suppress biofilm regrowth²⁴.

Once the microbial burden has been physically disrupted, it is significantly more susceptible to biocides and antibiotics while the biofilm works to reconstitute²⁵. Wound cleansing may be performed at this stage to remove surface contaminants, loose debris, slough, softened necrosis, bacteria and

Table 2 Considerations for diagnosing critical colonisation and localised infection in wound aetiologies (photos courtesy T Swanson)

Wound aetiology	Tips for diagnosis	Tips for management
Leg ulcer 	<p>Can be distinguished from chronic inflammation by being unilateral, warm to touch, increase in pain, and increased size of wound despite use of a therapeutic level of compression</p> <p>At least 3 of these 5 clinical signs of local infection:</p> <ul style="list-style-type: none"> ▪ Pain between dressing changes ▪ Periwound skin erythema ▪ Oedema ▪ Malodour ▪ High levels of exudate⁵ 	<p>Topical antimicrobial agents should be used with standard management practices such as graduated compression therapy</p> <p>Manage the increased exudate with an absorbent dressing suitable for use under compression therapy</p>
Diabetic foot ulcer 	<p>Signs and symptoms may be subtle (and up to half of infected DFUs will show no signs and symptoms), so it is important to be proactive in exploring if the wound is not progressing, particularly if 2 or more of these are present:</p> <ul style="list-style-type: none"> ▪ Local swelling or induration ▪ Erythema ▪ Local tenderness or pain ▪ Local warmth ▪ Presence of/increased exudate or purulent discharge^{14,15} <p>Probe to bone or exposed bone is likely indicative of osteomyelitis¹⁴</p>	<p>Listen to the patient regarding his/her subjective symptoms</p> <p>Manage factors such glycaemic control and offloading in consultation with a specialist (e.g. podiatrist)</p> <p>Use appropriate topical antimicrobial and/or systemic antibiotics for prevention and treatment per local protocol¹⁶, in consultation with a specialist</p>
Pressure ulcer 	<p>Local infection is likely in the presence of:</p> <ul style="list-style-type: none"> ▪ Lack of healing for 2 weeks ▪ Friable granulation tissue ▪ Malodour ▪ Increased pain in wound ▪ Increased periwound warmth ▪ Negative change in the nature of the wound drainage ▪ Increased necrotic tissue in the wound bed ▪ Pocketing or bridging of the wound bed ▪ No improvement/deterioration despite implementation of appropriate relief of pressure, shear and friction 	<p>Manage factors such as nutrition and pressure reduction/relief</p> <p>Reduce bacterial load and biofilm through cleansing, debridement and appropriate antimicrobial dressings¹⁷</p>
Surgical site wound 	<p>Post-operative symptoms include:</p> <ul style="list-style-type: none"> ▪ Redness and pain around the surgical area ▪ Drainage of cloudy fluid from the surgical wound ▪ Fever¹⁸ ▪ Wound breakdown/dehiscence 	<p>Identify risk factors before elective surgery and manage per local protocol</p> <p>Use an appropriate antimicrobial dressing or device (e.g. negative pressure wound therapy) to manage symptoms, exudate and bacterial burden¹⁹</p>
Superficial partial-thickness burn 	<p>It is important to differentiate between burn wound erythema (a normal process characterised by painless, blanchable redness <2cm from the burn edge for 3–5 days post-injury) and cellulitis, which presents with:</p> <ul style="list-style-type: none"> ▪ Advancing redness ▪ Warmth ▪ Tenderness²⁰ 	<p>Reassess within 24–72 hours of injury to monitor for increased depth (due to burn conversion) or infection</p> <p>Closely monitor for signs and symptoms of infection and response to treatment</p> <p>Use appropriate topical and/or systemic antimicrobial per local guidelines</p>

other microbes from the wound surface and surrounding skin²⁶. After debridement and cleansing, application of an antimicrobial dressing that is appropriate for the clinical indications (e.g. exudate and odour management), as well as safe for and acceptable to the patient, is recommended^{22,24,27,28} (Table 3). This is particularly important during the first 24 hours after debridement and cleansing, to protect the wound from the re-establishment of the microbial burden²⁹.

Role of antibiotics

Indiscriminate use of antimicrobials — antibiotics in particular — has made resistant organisms more prevalent³¹. During the next 50 years, microorganisms' drug resistance will increase, and new strains with resistance to a wide variety of agents will emerge, rendering many drugs ineffective³². In several countries that have launched large-

Table 3 Overview of topical antimicrobials³

Agent to reduce/prevent microbial burden	Additional rationales for use	Wound types	General guidance for use*
Enzyme alginate	<ul style="list-style-type: none"> Autolytic debridement Moisture balance Wound edge and epithelial cell protection 	<ul style="list-style-type: none"> PU DFU Acute wound Arterial ulcer Superficial partial-thickness burns 	<ul style="list-style-type: none"> Apply to wound and cover with a secondary dressing Can be used long-term due to no body absorption Contraindicated in patients with known sensitivity to alginate dressing or polyethylene glycol
Iodine (povidone, cadexomer)	<ul style="list-style-type: none"> Effective against MRSA³⁰ Reduced selection for bacterial resistance³⁰ 	<ul style="list-style-type: none"> VLU DFU Cavity wounds (cadexomer only) 	<ul style="list-style-type: none"> Use for 1 week, with dressing changes 2 to 3 times weekly If there are signs of improvement, continue use up to 2 weeks. If the wound does not improve after 1 week, discontinue use Contraindicated for long-term use, and in patients with known/suspected iodine sensitivity, and renal or thyroid diseases
Medical-grade honey	<ul style="list-style-type: none"> Autolytic debridement Odour management 	<ul style="list-style-type: none"> Leg ulcer Superficial or partial-thickness burn DFU PU Surgical wound Graft site 	<ul style="list-style-type: none"> Change dressing based on how quickly honey is diluted by exudate Ensure direct contact with the wound bed and use with a secondary dressing to manage exudate Monitor blood sugar levels in patients with diabetes Use with caution in patients with bee venom allergy Do not let the dressing dry out
Octenidine dihydrochloride	<ul style="list-style-type: none"> Autolytic debridement Donate moisture to the wound bed Wound cleansing 	<ul style="list-style-type: none"> Superficial or partial-thickness burn PU Leg ulcer DFU 	<ul style="list-style-type: none"> Apply directly to the wound bed and leave in contact for ≥5 minutes Can be used to soften dressings and loosen encrusted coatings before removal Contraindicated in patients with octenidine sensitivity, on exposed structures underlying the dermis
Polyhexamethylene biguanide (PHMB)	<ul style="list-style-type: none"> Odour management (dressing) Removal of encrusted dressings (solution) Debridement (gel) Wound bed preparation (gel and solution) Wound cleansing 	<ul style="list-style-type: none"> Partial-thickness burn Surgical wound Graft site Leg ulcer PU DFU 	<ul style="list-style-type: none"> Warm solution to body temperature, apply to gauze, then the wound, and cover and leave in contact for 10 minutes If using gel in dehydrated, deep, tunnelling or cavity wounds, first apply to a ribbon gauze Dressing can be left in place 5–7 days Contraindicated in patients with PHMB sensitivity Do not combine with other wound cleansers or ointments
Silver (metallic, nanocrystalline, ionic)	<ul style="list-style-type: none"> No known bacterial resistance in wounds Manage exudate, fill cavity wound, protect vulnerable tissue (combined with alginates or contact layers) 	<ul style="list-style-type: none"> All wound types With caution in children 	<ul style="list-style-type: none"> Use for 2 weeks. If there are signs of improvement, continue use up to 4 weeks. If there are no signs of improvement, discontinue use Do not use >4 weeks without strong clinical rationale Contraindicated for long-term use, over large surface areas and in patients with sensitivity to silver Use with caution in heavily exuding wounds (risk of maceration)
Silver sulfadiazine	<ul style="list-style-type: none"> Soothe painful wounds 	<ul style="list-style-type: none"> Partial- or full-thickness burn Leg ulcer PU 	<ul style="list-style-type: none"> Use for 1 week. If there are signs of improvement, use up to 2 weeks. If there are no signs of improvement, discontinue use Do not use >2 weeks Clean wound, and cover with 0.3–0.5cm thickness of cream and a secondary dressing Contraindicated in patients with sensitivity to silver sulfadiazine and sulfa drugs May result in the development of a false eschar

scale, coordinated efforts to reduce antibiotic usage, reports of incidences of resistant strains of bacteria, such as *Staphylococcus aureus* and *Clostridium difficile*, have decreased significantly³³.

It is therefore recommended that systemic antibiotics be used cautiously.

In addition, where possible, the choice of antibiotic prophylaxis should be matched to the organisms most likely to cause infection, using the local antibiotic formulary guidelines to ensure the most appropriate antibiotic, dose, timing of administration and duration of use³⁴. Antibiotic prophylaxis is indicated in

patients who have systemic risk factors that make infection likely (e.g. poor vascularity, compromised immune systems), particularly in wounds at high risk of becoming infected, such as contaminated wounds, wounds with large areas of necrotic tissue, and high-risk anatomical sites such as the sacrum,

Box 2 Potential triggers for systemic antibiotic use*

- Abnormal/absent granulation or necrosis
- Pocketing, tunnelling, maceration
- Static or enlarged wound size
- Erythema spreading >2cm around the wound
- Appearance of or changes in nature of pain
- Wound deepening to involve structures under the skin and subcutaneous tissues[†]
- Body temperature >37°C[†]
- Heart rate >90 beats per minute[†]

*Not exhaustive

[†]Systemic infection only

hand or foot (Box 1)^{31,35}. For example, if a DFU shows signs of critical colonisation or local infection, antibiotics should be initiated due to the potential for rapid onset of infection, as well as the risk of amputation if infection settles in¹³.

If there are signs of systemic infection, systemic antibiotics should be initiated; a topical antimicrobial can also be used if localised effect is desired, and the wound status allows dressing application and change without further damage to surrounding structures (Box 2). If the patient and wound are showing signs of spreading infection, use of systemic antibiotic therapy may be considered in addition to a topical antimicrobial (Box 2).

Monitoring progress

A multidisciplinary approach, coupled with a treatment pathway that enables timely referral to specialists, is important for optimal outcomes. Thorough, ongoing assessment should be employed to evaluate the progression of the wound (according to treatment goals) and the effect of the current

treatment on meeting these goals³⁶. The results of assessment should be clearly documented in the patient record and care plan, with any changes and a clear rationale for such changes recorded³⁶.

Outcomes can be measured by improvement of wound and periwound conditions, and quality-of-life indicators (e.g. pain, odour)³⁷. Healing trajectories are noted in the literature:

- **It is reported that most wounds should heal within about 4 weeks³⁷**
- **Burns healing 2% per day have positive indications for healing and patient survival³⁸**
- **DFUs that do not reduce in area by ≥50% within 4 weeks are unlikely to heal by 12 weeks³⁹**

The clinician should review the wound 1 week after presentation (or per local protocol), looking for positive indicators such as decreased pain, exudate (level and type), odour and oedema; and improved periwound skin (intact, decreased erythema and oedema), wound edges (intact, non-inflamed) and wound bed (increased granulation tissue, decreased non-viable tissue).

If these positive indicators and improvements are not being seen, it is important to investigate whether infection or an underlying condition is the cause. If antibiotics have been initiated, a positive indicator within 24–48 hours should be expected.

Frequency of dressing change should be based on regular, ongoing assessment,

dressing indications (e.g. exudate, odour, infection management), manufacturer instructions and patient factors (e.g. sensitivity to materials, concordance with dressing choice). Change treatment if dressing performance does not meet expectations for the clinician or patient, or if the patient experiences any adverse effects from treatment. The changing condition of the wound may also initiate a change in the care plan (e.g. if symptoms have resolved or if new symptoms present).

Where wound healing is not an achievable outcome, it is as important to manage the wound locally as it is to implement holistic assessment and management, including nutrition, stress (physical and emotional/psychological), lifestyle factors, quality of life and medication, in concordance with the wishes of the patient and their families.

Supported by an unrestricted educational grant from Acelyt. The views expressed in this Made Easy do not necessarily reflect those of Acelyt.

Author details

Swanson T¹, Grothier L², Schultz G³

1. Chair International Wound Infection Institute, Nurse Practitioner Wound Management, South West Healthcare, Warrnambool, Australia;
2. Consultant Nurse Tissue Viability/Service Manager, Tissue Viability Centre, Provide CIC, St Peter's Hospital, Maldon, UK
3. Professor, Department of Obstetrics and Gynecology, Institute for Wound Research, University of Florida, Gainesville, USA

Summary

Wounds that have been clinically diagnosed as infected could be treated with a topical antimicrobial that is appropriate for the clinical indications (e.g. exudate and odour management) as well as safe for and acceptable to the patient. Systemic antibiotics should be considered and used cautiously and in consultation with a specialist member of the multidisciplinary team. A multidisciplinary approach, coupled with a treatment pathway that enables timely referral to specialists, is important for optimal outcomes. Thorough, ongoing assessment should be employed to evaluate the progression of the wound (according to the treatment goals) and the effect of treatment on meeting treatment goals.

To cite this publication

Swanson T, Grothier L, Schultz G. Wound Infection Made Easy. *Wounds International* 2014. Available from: www.woundsinternational.com

References

1. Thomas B, Bishop J. *The Manual of Dietetic Practice* (4th edn). Blackwell Publishing: London, 2007.
2. Penhallow K. A review of studies that examine the impact of infection on the normal wound-healing process. *J Wound Care* 2005;14(3):123–6.
3. Wounds UK *Best Practice Statement. The use of topical antimicrobial agents in wound management* (3rd edn). London: Wounds UK, 2013. Accessed 08.10.2014 at: <http://www.wounds-uk.com/best-practice-statements/best-practice-statement-the-use-of-topical-antimicrobial-agents-in-wound-management>
4. Moore K, Huddleston E, Stacey MC, Harding KG. Venous leg ulcers — the search for a prognostic indicator. *Int Wound J* 2007;4(2):163–72.
5. World Union of Wound Healing Societies. *Principles of best practice: Wound infection in clinical practice. An international consensus*. London: MEP Ltd, 2008. Accessed 08.10.2014 at <http://www.woundsinternational.com/clinical-guidelines/wound-infection-in-clinical-practice-an-international-consensus>
6. Leaper D. Silver dressings: their role in wound management. *Int Wound J* 2006;3(4):282–94.
7. Levine NS, Lindberg RB, Mason AD, Pruitt BA Jr. The quantitative swab culture and smear: A quick, simple method for determining the number of viable aerobic bacteria on open wounds. *J Trauma* 1976;16(2):89–94.
8. Gardner SE, Frantz RA, Saltzman CL, et al. Diagnostic validity of three swab techniques in identifying chronic wound infection. *Wound Repair Regen* 2006;14(5):548–57.
9. Angel DE, Lloyd P, Carville K, Santamaria N. The clinical efficacy of two semi-quantitative wound-swabbing techniques in identifying the causative organism (s) in infected cutaneous wounds. *Int Wound J* 2011;8(2):176–185.
10. Cutting K, Harding K. Criteria for identifying wound infection. *J Wound Care* 1994;3(4):198–201.
11. Cutting K, White RJ, Maloney P, Harding KG. Clinical identification of wound infection: a Delphi approach. In: *European Wound Management Association Position Document. Identifying criteria for wound infection*. London: MEP Ltd, 2005.
12. Gardner SE, Frantz RA, Doebbeling BN. The validity of the clinical signs and symptoms used to identify localized chronic wound infection. *Wound Repair Regen* 2001;9(3):178–186.
13. Wounds International. *International Best Practice Guidelines: wound management in diabetic foot ulcers*. London: Wounds International, 2013. Accessed 10.10.2014 at: <http://www.woundsinternational.com/clinical-guidelines/best-practice-guidelines-wound-management-in-diabetic-foot-ulcers>
14. Lipsky BA, Berendt AR, Cornia PB, et al. 2012 Infectious Diseases Society of America clinical practice guideline for the diagnosis and treatment of diabetic foot infections. *Clin Infect Dis* 2012;54(12):132–173.
15. Lavery LA, Armstrong DG, Wunderlich RP, et al. Risk factors for foot infections in individuals with diabetes. *Diabetes Care* 2006;29(6):1288–93.
16. Lipsky BA, Hoey C, Cruciani M, Mengoli C. Topical antimicrobial agents for preventing and treating foot infections in people with diabetes (Protocol). *Cochrane Database of Systemic Reviews* 2014;3:1–22.
17. National Pressure Ulcer Advisory Panel, European Pressure Ulcer Advisory Panel, Pan Pacific Pressure Injury Alliance. *Prevention and Treatment of Pressure Ulcers: Clinical Practice Guideline*. Emily Haesler (Ed). Cambridge Media: Perth, Australia; 2014.
18. Centers for Disease Control and Prevention (CDC). Frequently asked questions about surgical site infections. 2012. Accessed 20.10.2014 at: http://www.cdc.gov/HAI/ssi/faq_ssi.html
19. Keast D, Swanson T, Sussman G, et al. Ten top tips: managing surgical site infections. *Wounds International* 2014;5(3):13–8. Accessed 20.10.2014 at: <http://www.woundsinternational.com/practice-development/ten-top-tips-managing-surgical-site-infections>
20. Mitchell KB, Leahy NE, Gallagher JJ. Treating infection in burns. *Contemporary Pediatr* 2013;30(9):26–32.
21. James GA, Swogger E, Wolcott R, et al. Biofilms in chronic wounds. *Wound Repair Regen* 2008;16(1):37–44.
22. Wolcott RD, Rhoads DD. A study of biofilm-based wound management in subjects with critical limb ischaemia. *J Wound Care* 2008;17(4):145–55.
23. Phillips PL, Wolcott RD, Fletcher J, Schultz GS. Biofilms Made Easy. *Wounds International* 2010. Accessed 20.10.2014 at: <http://www.woundsinternational.com/made-easys/biofilms-made-easy>
24. Schultz G. Understanding biofilm-based wound care: What you need to know. *Wounds International* webcast. 2011. Accessed 25.09.2014 at <http://www.woundsinternational.com/webcasts/understanding-biofilm-based-wound-care-what-you-need-to-know>
25. Wolcott RD, Rumbaugh KP, James G, et al. Biofilm maturity studies indicate sharp debridement opens a time-dependent therapeutic window. *J Wound Care* 2010;19(8):320–8.
26. Rodeheaver RT, Ratliff CR. Wound cleansing, wound irrigation and wound disinfection. In: Rodeheaver GT, Krasner DI, Sibbald RG eds. *Chronic wound care: A clinical source book for healthcare professionals*. Malvern, USA: HMP Communications, 2007.
27. Wolcott RD, Rhoads DD, Dowd SE. Biofilms and chronic wound inflammation. *J Wound Care* 2008;17(8):333–41.
28. Fonseca A. Biofilms in wounds: An unsolved problem? *EWMA J* 2011;11(2):10–2.
29. Wound Healing Management Node Group Evidence summary: wound infection: biofilms defined and described. *Wound Pract Res* 2012;20(4):187–9.
30. Mertz PM, Oliveira-Gandia MF, Davis SC. The evaluation of cadexomer iodine dressing on methicillin-resistant *Staphylococcus aureus* (MRSA) in acute wounds. *Dermatol Surg* 1999;25(2):89–93.
31. European Wound Management Association (EWMA). EWMA document. Antimicrobials and non-healing wounds: evidence, controversies and suggestions. *J Wound Care* 2013;22(Suppl 5):S1–S89.
32. Davies SC. Annual report of the chief medical officer. Volume two, 2011. *Infections and the rise of antimicrobial resistance*. London: Department of Health, 2013. Accessed 09.10.2014 at: <https://www.gov.uk/government/publications/chief-medical-officer-annual-report-volume-2>
33. Pittet D. The evolving threat of antimicrobial resistance: options for action. Presented 8 March 2012 at: Book launch event, WHO Headquarters, Geneva. Accessed 10.10.2014 at: <http://www.who.int/patientsafety/implementation/amr/en/>
34. National Institute for Health and Care Excellence (NICE). *Surgical site infection. NICE quality standard 49: surgical site infection*. London: NICE, 2013. Accessed 10.10.2014 at: guidance.nice.org.uk/qs49
35. World Health Organization. Prevention and management of wound infection. Accessed 09.10.2014 at: http://www.who.int/hac/techguidance/tools/guidelines_prevention_and_management_wound_infection.pdf
36. Ousey K, Atkin L. *Optimising the patient journey Made Easy*. London: Wounds UK, 2013. Accessed 10.10.2014 at: <http://www.wounds-uk.com/made-easy/optimising-the-patient-journey-made-easy>
37. Bernatchez SF, Menon V, Stoffel J, et al. Nitric oxide levels in wound fluid may reflect the healing trajectory. *Wound Repair Regen* 2013;21(3):410–7.
38. Nitzschke S, Aden J, Serio-Melvin M, et al. Wound healing trajectories in burn patients and their impact on mortality. *J Burn Care Res* 2014 Aug 20. [Epub ahead of print].
39. Warriner RA, Snyder RJ, Cardinal MH. Differentiating diabetic foot ulcers that are unlikely to heal by 12 weeks following achieving 50% percent area reduction at 4 weeks. *Int Wound J* 2011;8(6): 632–637.